

CLASSIFICATION UNCLASSIFIED	DOCUMENT IDENTIFICATION NO. BNWL-CC-1850
BATTELLE NORTHWEST BATTELLE MEMORIAL INSTITUTE PACIFIC NORTHWEST LABORATORY POST OFFICE BOX 999 / RICHLAND, WASHINGTON 99352	COPY AND SERIES NO.
	DATE September, 1968
TITLE AND AUTHOR <u>ENVIRONMENTAL STATUS OF THE HANFORD RESERVATION FOR JANUARY-JUNE, 1968</u>	CONTRACT PROJECT NO. <input checked="" type="checkbox"/> - 1830 <input type="checkbox"/> - 1831
By Environmental Evaluation Radiation Protection TECHNICAL SERVICES DIVISION	RESERVED FOR TECH. INFO. USE

BEST AVAILABLE COPY

D I S T R I B U T I O N

NAME	COMPANY	LOCATION	NAME	COMPANY	LOCATION
<u>Battelle-Northwest</u>			<u>ARHCO, Inc.</u>		
		<u>Bldg.</u> <u>Area</u>			<u>Bldg.</u> <u>Area</u>
1.	D. W. Alton	3705 300	33.	G. E. Backman	222 T 200 W
2.	P. J. Blumer	3705 300	34.	D. J. Brown	271 U 200 W
3.	L. A. Carter	328 300	35.	G. L. Hanson	271 B 200 E
4.	J. P. Corley	3705 300	36.	B. J. McMurray	202 S 200 W
5.	T. H. Essig	3705 300	37.	Files	
6.	W. L. Fisher	3701-L 300	<u>Atomic Energy Commission - Richland Operations Office</u>		
7.	R. F. Foster	326 300	38.	A. Brunstad	Fed. 700
8.	R. B. Hall	328 300	39.	P. A. Craig	Fed. 700
9.	J. F. Honstead	3702 300	40.	W. Devine, Jr.	Fed. 700
10.	W. C. Horton	3705 300	41.	W. E. Lotz	Fed. 700
11.	R. T. Jaske	713 700	42.	C. L. Robinson	Fed. 700
12.	J. J. Jech	713 700	43.	R. B. St. John	Fed. 700
13.	H. V. Larson	3705 300	44.	M. W. Tiernan	Fed. 700
14.	J. M. Selby	328 300	45.	Technical Library	
15.	J. K. Soldat	3702 300	<u>Douglas-United Nuclear, Inc.</u>		
16.	W. L. Templeton	1704 100 F	46.	D. A. Baker	1100 N 100 N
17.	F. W. Woodfield	703 700			
18-19.	C. B. Wooldridge	3705 300			
<u>20-22. Technical Information</u>					
<u>23-32. Extra Copies</u>					

ROUTE TO	PAYROLL NO.	COMPANY	LOCATION	FILES ROUTE DATE	SIGNATURE AND DATE
<i>J. R. Young</i>	33539		Math	JUL 14 1971	
<i>Please return to me</i>			<i>Math/300</i>		
<i>J. R. Young</i>	33539		<i>Math</i>	FEB 5 1974	

THIS DOCUMENT IS PUBLICLY AVAILABLE

INFORMATION CONCERNING USE OF THIS REPORT

PATENT STATUS

This document copy, since it is transmitted in advance of patent clearance, is made available in confidence solely for use in performance of work under contracts with the U. S. Atomic Energy Commission. This document is not to be published nor its contents otherwise disseminated or used for purposes other than specified above before patent approval for such release or use has been secured, upon request, from the Chief, Chicago Patent Group, U. S. Atomic Energy Commission, 9800 So. Cass Ave., Argonne, Illinois.

PRELIMINARY REPORT

This report contains information of a preliminary nature prepared in the course of work under Atomic Energy Commission Contract AT(45-1)-1830. This information is subject to correction or modification upon the collection and evaluation of additional data.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

PACIFIC NORTHWEST LABORATORY

RICHLAND, WASHINGTON

operated by

BATTELLE MEMORIAL INSTITUTE

for the

UNITED STATES ATOMIC ENERGY COMMISSION UNDER CONTRACT AT(45-1)-1830

UNCLASSIFIED

CLASSIFICATION

DOCUMENT IDENTIFICATION NO.

BNWL-CC-1850

BATTELLE **B** NORTHWEST
BATTELLE MEMORIAL INSTITUTE PACIFIC NORTHWEST LABORATORY
POST OFFICE BOX 999 / RICHLAND, WASHINGTON 99352

COPY AND SERIES NO.

DATE

September, 1968

TITLE AND AUTHOR

CONTRACT

PROJECT NO.

- 1830
 - 1831

RESERVED FOR TECH. INFO. USE

D I S T R I B U T I O N

NAME	COMPANY	LOCATION	NAME	COMPANY	LOCATION
<u>Douglas-United Nuclear, Inc. (Cont.)</u>					
		<u>Bldg.</u>	<u>Area</u>		
47.	P. A. Carlson	1760 D	100		
48-49.	C. D. Corbit	1760 H	100 H		
50.	R. G. Geier	1760 D	100		
51.	W. S. Nechodom	1101 N	100 N		
52.	F. E. Owen	1760 H	100 H		
53.	L. F. Reilly	3717	300		
54.	Files				
<u>J. A. Jones Construction Company</u>					
55.	L. C. Roos	1256	3000		

ROUTE TO	PAYROLL NO.	COMPANY	LOCATION	FILES ROUTE DATE	SIGNATURE AND DATE

UNCLASSIFIED

CLASSIFICATION

ENVIRONMENTAL STATUS OF THE
HANFORD RESERVATION FOR JANUARY- JUNE, 1968

By

Environmental Evaluations Staff
Radiation Protection Department
TECHNICAL SERVICES DIVISION

edited by C. B. Wooldridge

September, 1968

UNCLASSIFIED

TABLE OF CONTENTS

	<u>Page</u>
I. Introduction	1
II. Surveillance Summary	2
III. Columbia River Water	4
A. Raw Water	4
B. Drinking Water	7
IV. Swamps, Ditches, and Ponds	10
A. Waste Water	10
B. Game Birds	19
V. The Atmosphere	20
A. Iodine-131 and Total Beta Activity	20
B. Total Alpha Activity	30
VI. Radiation Surveys	35
A. Surface Contamination	35
1. Hanford Roads	35
2. Control Plots	38
3. Waste Disposal Sites	40
4. Aerial Survey	42
B. External Radiation Exposure Rates	43
1. On Plant	43
2. At the Columbia River Shoreline	52
3. Below the Surface of the Columbia River	58

LIST OF FIGURES

		<u>Page</u>
1	Radioactivity of Columbia River Raw Water (Grab) Samples	6
2-3	Radioactivity of Drinking Water (Grab) Samples	8-9
4-8	Radioactivity of Waste Water Samples	14-18
9-15	Iodine-131 and Total Beta Activity in the Atmosphere	23-29
16-19	Total Alpha Activity in the Atmosphere	31-34
20-26	External Radiation On-Plant	45-51
27-28	External Radiation at the Columbia River Shoreline	55-56
29-30	External Radiation Below the Surface of the Columbia River	60-61

LIST OF MAPS

		<u>Page</u>
1	Drinking and Raw Water Sampling Locations	5
2	Swamps, Ditches, and Pond Water Sampling Locations	13
3	Air Sampling Stations	22
4	Hanford Road Surveys	37
5	Control Plot Locations	39
6	Underground and Surface Radioactivity	41
7	On-Plant External Radiation Exposure Rate Measurement Points	44
8	Shoreline Survey Locations	54
9	Exposure Rate Measurement Points in the Columbia River	59

LIST OF TABLES

	<u>Page</u>
1 Radionuclide Concentrations in the Columbia River at Priest Rapids Dam (Jan.-Mar.) and Priest Rapids Gauge Station (April-Present)	4
2 Radionuclide Concentrations in Waste Water Samples	10
3 231-Z Waste Pit	11
4 300 Area Leach Trench and River Shoreline Seepage Area	12
5 300 Area Process Pond - Fluoride Concentrations	12
6 Atmospheric ^{131}I and Total Beta Concentrations - Averages	21
7 Atmospheric Total Alpha Concentrations - Averages	30
8 Average Exposure Rates	43
9 Monthly River Shoreline Surveys - January-June, 1968 (Maximum Readings)	57

ENVIRONMENTAL STATUS OF THE
HANFORD RESERVATION FOR JANUARY-JUNE, 1968

I. INTRODUCTION

This report summarizes data collected during the first six months of 1968 from locations within the Hanford plant boundaries for the routine environmental surveillance program, under the direction of the Environmental Evaluations staff. These environmental data are reported here for the information of the Richland Operations Office of the Atomic Energy Commission and its contractors.

To show long-term trends and facilitate comparisons, the graphs in this report show 14 months of data - for the current half-year and the preceeding 8 months. However, the reader may also wish to refer to BNWL-CC-1197-6, "Environmental Status of the Hanford Reservation for November-December, 1967 (Annual Summary)" which contains a complete summary of 1967 data. Ground water data are not included in this report but will be presented in a BNWL report, "Radiological Status of the Ground Water Beneath the Hanford Project, January-June, 1968".

Data from off-site sampling locations is presented in the series of monthly reports, BNWL-778-___, "Environmental Surveillance in the Vicinity of Hanford for . . . , 1968".

The term "analytical limit", as used herein, is defined as the concentration at which the laboratory can measure a radionuclide with a precision of ± 100 per cent at the 90 per cent confidence level. The detection limit for a specific radionuclide varies with sample type, sample size, counting time, and the amounts of interfering radionuclides present. The "analytical limits" were chosen to represent upper bounds to these fluctuating detection limits.

The majority of the data presented in this report was supplied by the U. S. Testing Co., Inc., which performed all routine radioassays of environmental samples.

II. SURVEILLANCE SUMMARY - JANUARY-JUNE, 1968

A summary of surveillance highlights for the period is given below with references to the page number(s) where more detail on the subject appears.

Liquid Waste Disposal

Three production reactors remained operational following the retirement of "B" reactor in mid-February. Dose rates and surface contamination measurements at shoreline locations downstream from the reactors, including Ringold, the Power Line Crossing, and Richland, showed significant increases between October of 1967 and May of 1968. Dose rate increases of up to 50% accompanied increased prevalence of particulates trapped in debris or in the rocks. The maximum shoreline exposure rate during the first six months of 1968, 1500 R/hr, was measured below 100-K. Particles found on the shoreline just below the plant boundary read as much as 75,000 c/m (GM). At the farthest downstream measurement location, sacajawea, patches of foam measuring 9,000 c/m (GM) were detected in May. (See p. 52).

Disposal of low-level radioactive wastes to open swamps, ponds, and trenches resulted in several unusual events. Total beta concentrations exceeded the limit for open waters (50,000 pCi/l, Ref. AECM 0510) in T Swamp during May and June; the principal gamma emitters present were ^{137}Cs and ^{134}Cs . (See p. 10). Releases through the Purex Chemical Sewer resulted in several instances of high ^{131}I concentrations in B Swamp with a maximum of 47,000 pCi ^{131}I /l occurring in April (p. 11). Total alpha concentrations temporarily exceeded 50,000 pCi/l at the 200 West 222-S Swamp in April (p. 10). In addition, total alpha concentration in samples from the 231 Z Waste Pit exceeded the limit of 50,000 pCi/l in February, with the maximum alpha activity of 420,000 pCi/l in a sample collected February 23. The waste pit was returned on March 21, 1968.

Gaseous Waste Disposal

During the first six months of 1968, several transient increases in total beta, total alpha, and ^{131}I concentrations were observed. A general increase in the routine beta concentrations at most sample locations was attributed to fallout. (See p. 20-33).

In March, a particle filter from 200 West Redox indicating $3.4 \text{ pCi}\beta/\text{m}^3$ and $0.91 \text{ pCi}\alpha/\text{m}^3$ for the period 3-11 to 3-13 was analyzed by alpha spectrometry. The activity was radon-thoron and their respective daughters.

Other transient increases in atmospheric beta activity occurred at 100-N and 200 East Main Gate in early January, at 200-East East Center in early February and at 100-K in late May, with a maximum concentration of $2.5 \text{ pCi}\beta/\text{m}^3$ at 200-East East Center.

Gaseous Waste Disposal (Continued)

Temporary increases in atmospheric ^{131}I concentrations were also observed at 100-F, 100-B, and 200-East East Center in March and April, with a maximum concentration of $0.24 \text{ pCi}^{131}\text{I}/\text{m}^3$ at 100-B.

Increases in ^{131}I at the 300 Area in January and March and at the 700 Area in March resulted from temporary increases in ^{131}I emission rates from laboratories in the 300 Area. The maximum concentration ($0.54 \text{ pCi}^{131}\text{I}/\text{m}^3$) was noted in March at the 300 Area.

In early June, alpha spectrometry of the higher than usual activity ($0.05 \text{ pCi}/\text{m}^3$) on a 300 Area filter yielded the following results: $^{234}\text{-}^{235}\text{U}$ - 42%, ^{232}Th - 36%, ^{238}U - 20%, and ^{239}Pu - 1.8%.

Surface Contamination

Routine surveys of plant roads with a gamma spectrometer during the first six months of 1968 detected a total of 10 instances of particulate deposition, as compared to a total of seven such occurrences for all of 1967. Most of the increased road deposition appeared to be associated with radioactive material transport in the vicinity of the 100 Areas. Generally, contamination was either attached to the road surface or embedded in the sand and gravel beside the road. Follow-up surveys of adjacent off-road locations showed no additional contamination. Radiation levels of these particulates were as high as $100,000 \text{ c/m (GM)}$ and $150 \text{ mrads/hr (Surface-Juno)}$. (See p. 52).

No deposition of radioactive particulates on the routine control plots was detected during January-June, 1968.

Several retired waste burial grounds were surveyed and the appropriate custodians were notified of conditions at 100-F, the Wye, and the 300 Area North burial grounds which required attention. Radiation levels to $75,000 \text{ c/m (GM)}$ were observed (p. 40).

Fallout from Nuclear Weapons Testing

In early January, atmospheric beta activity increases were attributed to fallout from an announced nuclear weapons test of December 24, 1967. An influx of fallout in May and June resulted in higher than normal beta concentrations. With these influxes of fallout, corresponding increases in external exposure rates were noted (p. 43).

III. COLUMBIA RIVER WATER

The locations from which raw and drinking water samples were obtained are shown in Map 1.

A. Raw Water


Weekly cumulative samples of raw Columbia River water from Priest Rapids Dam were analyzed for total alpha and total beta. In addition, a monthly composite was analyzed for ^3H , ^{90}Sr , $^{95}\text{Zr-Nb}$, and ^{106}Ru . Results appear in Table I. The numbers in parentheses are the analytical limits for each analysis.

TABLE I
 RADIONUCLIDE CONCENTRATIONS IN THE COLUMBIA RIVER
 AT PRIEST RAPIDS DAM (JAN.-MAR.) AND
 PRIEST RAPIDS GAUGE STATION (APRIL-JUNE)


<u>Month</u>	<u>^3H (pCi/l)</u>	<u>^{90}Sr (pCi/l)</u>	<u>Total α (pCi/l)</u>	<u>Total β (c/m/ml)</u>
January	2300 (1000)	0.53 (0.50)	<1.2 (1.0)	<0.005
February	2500	<0.50	<1.1	<0.005
March	1500	<0.50	<1.0	<0.005
April	<1000	<0.50	<1.0	<0.006
May	*	*	<1.2	<0.005
June	*	<0.50	<1.0	<0.005

* No analysis made.

The total beta activity in river water at Ringold was obtained from analyses of weekly grab samples. Results of these analyses appear in Figure 1.




RADIOACTIVITY OF COLUMBIA RIVER RAW WATER (GRAB) SAMPLES




B. Drinking Water

Total beta analyses of drinking water samples taken from 100-K, 100-N, 200-W, and the B-Y Telephone exchange are shown in Figures 2 and 3. The 100-K sampling location was added on January 1, 1968. As expected, total beta concentrations at 100-K have been significantly lower than those observed at 100-N. The 100-N and 200-W locations were switched to a weekly sampling frequency on January 1, 1968, in order to more closely follow the short-term changes in total beta concentrations. The monthly GI tract dose shown in Figure 3 is estimated from isotopic analyses of drinking water samples collected at 100-N. This estimate is based on an assumed intake of 1.2 liters per day, 5 days per week.

RADIOACTIVITY OF DRINKING WATER (GRAB) SAMPLES



RADIOACTIVITY OF DRINKING WATER (GRAB) SAMPLES



IV. SWAMPS, DITCHES, AND PONDS

A. Water

Open waters which may be used by migratory waterfowl are routinely sampled at the locations shown in Map 2. The sampling frequencies for all locations were changed to monthly on January 1, 1968. Figures 4-6 show data collected from swamps and ditches located within or near the 200 West Area. Figure 7 shows data collected from swamps and ditches located within and near the 200 East Area, and Figure 8 shows data collected from the 300 Area pond.

All swamp samples collected between January and June were quantitatively analyzed for significant radionuclides. These results appear in Table 2. Increases in radionuclide concentrations beyond the expected range of variation were noted at the following locations during January-June: U Swamp North (beta), T Swamp North (beta), 222-S Swamp (alpha), Gable Swamp North (alpha) and Purex Chemical Sewer (alpha).

TABLE 2

RADIONUCLIDE CONCENTRATIONS IN WASTE WATER SAMPLES

(pCi/l)

	<u>Date</u>	<u>⁵¹Cr</u>	<u>⁹⁵Zr-Nb</u>	<u>¹⁰³Ru</u>	<u>¹⁰⁶Ru</u>	<u>¹³⁴Cs</u>	<u>¹³⁷Cs</u>	<u>¹⁴⁴Ce-Pr</u>	<u>²³⁹Np</u>
<u>T Swamp North</u>	1-19	640	ND*	ND	<440	ND	960	<530	ND
	3-22	1100	ND	ND	<440	660	2300	<530	ND
	4-19	2000	ND	<57	2700	<150	430	ND	ND
	5-24	750	<52	ND	ND	930	3700	ND	ND
	6-21	<470	ND	ND	ND	240	720	ND	200
	<u>Date</u>	<u>⁵¹Cr</u>	<u>¹³¹I</u>	<u>¹³⁷Cs</u>	<u>¹⁴⁴Ce-Pr</u>				
<u>U Swamp North</u>	1-19	1900	ND	68	ND				
	2-23	2200	ND	90	<520				
	3-22	1400	ND	79	ND				
	4-19	1300	ND	55	<520				
	5-24	1100	ND	160	<530				
6-21	700	<49	350	ND					
	<u>Date</u>	<u>⁵¹Cr</u>	<u>⁶⁵Zn</u>	<u>¹³⁷Cs</u>	<u>¹⁴⁴Ce-Pr</u>				
<u>222-S Swamp</u>	1-19	1400	<96	<51	ND				
	2-23	1600	ND	<51	<520				
	3-22	1700	ND	<51	ND				
	4-19	1300	ND	<51	<520				
	5-24	1000	ND	<51	ND				
6-21	<480	<96	<51	ND					

* ND - Not detected.

A. Water (Continued)

TABLE 2 (Continued)

	<u>Date</u>	<u>51Cr</u>	<u>65Zn</u>	<u>137Cs</u>	<u>144Ce-Pr</u>		
Redox	1-19	2500	<96	<51	ND		
<u>North East</u>	2-23	2200	ND	<51	ND		
	3-22	2000	ND	<51	ND		
	4-19	1700	ND	ND	<520		
	5-24	880	<96	<51	ND		
	6-21	<480	ND	<51	ND		
	<u>Date</u>	<u>51Cr</u>	<u>103Ru</u>	<u>106Ru</u>	<u>131I</u>	<u>137Cs</u>	<u>144Ce-Pr</u>
Purex Chemi- <u>cal Sewer</u>	1-19	4300	160	6200	ND	<51	ND
	2-23	3500	ND	<440	ND	<51	<520
	3-22	ND	ND	ND	2,800	<51	ND
	4-19	ND	ND	ND	72,000	<51	ND
	5-24	1300	ND	ND	ND	<51	<520
	6-21	ND	ND	ND	64,000	<51	ND
	<u>Date</u>	<u>51Cr</u>	<u>65Zn</u>	<u>106Ru</u>	<u>131I</u>	<u>137Cs</u>	<u>144Ce-Pr</u>
B Swamp	1-19	150	ND	1400	ND	<51	<530
<u>North</u>	2-23	150	<96	<440	ND	ND	ND
	3-22	ND	ND	ND	860	<51	ND
	4-19	ND	ND	ND	47,000	<51	ND
	5-24	ND	ND	ND	790	<51	ND
	6-21	ND	ND	ND	3,100	<51	ND
	<u>Date</u>	<u>51Cr</u>	<u>65Zn</u>	<u>137Cs</u>	<u>144Ce-Pr</u>		
Gable Swamp	1-19	2100	<96	56	<1200		
<u>North</u>	2-23	2000	ND	280	<520		
	3-22	1600	ND	100	ND		
	4-19	1400	ND	150	<520		
	5-24	740	<96	96	ND		
	6-21	<480	<96	86	ND		

Results for the 231-Z Waste Pit (a temporary sampling location) are shown below. The waste pit was retired on March 21, 1968.

TABLE 3

231-Z WASTE PIT

<u>Date</u>	<u>Total Alpha</u> (pCi/l)	<u>Total Beta</u> (pCi/l)
1-5-68	1,300	270
1-12-68	220	210
1-19-68	500	330
1-26-68	1,100	95

A. Water (Continued)

TABLE 3 (Continued)

<u>Date</u>	<u>Total Alpha (pCi/l)</u>	<u>Total Beta (pCi/l)</u>
2-2-68	220,000	5,100
2-9-68	3,600	320
2-16-68	6,200	550
2-23-68	420,000	7,600
3-1-68	19,000	930
3-8-68	12,000	580
3-15-68	27,000	1,100

300 Area Leach Trench - Samples were periodically collected from the 300 Area sanitary waste disposal site for coliform, enterococci (a fecal organism) and BOD (biochemical oxygen demand) analyses.

TABLE 4

<u>Date</u>	<u>300 Area Leach Trench</u>		
	<u>Coliform/100 ml</u>	<u>Enterococci/100 ml</u>	<u>BOD mg/l</u>
3-19	180,000	20,000	-
6-18	200,000	350,000	18


<u>Date</u>	<u>River Shoreline Seepage Area</u>		
	<u>Coliform/100 ml</u>	<u>Enterococci/100 ml</u>	<u>BOD mg/l</u>
3-19	38	31	-
6-18	32	59	1.5

300 Area Process Pond - Weekly cumulative samples were collected from the 300 Area process pond throughout this report period. Results for total beta, uranium, nitrate, and hexavalent chromium appear in Figure 8. Monthly averages of the fluoride concentrations appear below.

TABLE 5


300 AREA PROCESS POND - FLUORIDE CONCENTRATIONS

<u>Month</u>	<u>Average Concentrations (ppm)</u>
January	3.7
February	3.0
March	2.5
April	3.3
May	2.7
June	3.5



SWAMPS, DITCHES, AND POND WATER SAMPLING LOCATIONS


● Sampling Location




RADIOACTIVITY OF WASTE WATER SAMPLES

△ Alpha

● Beta





RADIOACTIVITY OF WASTE WATER SAMPLES




RADIOACTIVITY OF WASTE WATER SAMPLES

△ Alpha

● Beta




RADIOACTIVITY OF WASTE WATER SAMPLES




WASTE WATER SAMPLE ANALYSES - 300 AREA


PROCESS POND INLET



PROCESS POND INLET



PROCESS POND INLET



IV. SWAMPS, DITCHES, AND PONDS (Continued)

B. Game Birds

No game bird samples were collected near liquid waste disposal sites during this report period.

V. THE ATMOSPHERE

A. Iodine-131 and Total Beta Activity

Results of routine sampling of the atmosphere at 19 locations within the Hanford Reservation (Map 3) are shown in Figures 9-15. At most locations the sampling equipment was contained within a small building designated "614". The sample is taken at a flow rate of 1.5 cfm through HV-70 filter paper and then passed through a solution of NaOH for ^{131}I collection. The normal sampling period is one week. "Total Beta" represents the activity of particulates collected on filter paper during the sampling period. During the first six months of 1968 several transient increases in total beta and/or ^{131}I concentrations were observed, as well as a general increase in the routine beta concentration at most sample locations. This increase was attributed to fallout. A month-by-month summary of each change in concentration follows.

January-February - Increased atmospheric beta activity starting early in January was attributed to fallout following an announced nuclear weapons test on December 24, 1967.

In January, increases were noted in ^{131}I concentrations at the 300 Area following above normal ^{131}I emissions from laboratory buildings.

Increased beta concentrations were noted at 100-N and 200-East Main Gate in early January and 200-East East Center in early February but could not be explained on the basis of known plant releases.

Air sampling was initiated at 200-West Northeast during the latter part of February.

March-April - A particle filter removed after running from 3-11 to 3-13 at 200-West Redox indicated a beta concentration of 3.4 pCi/m^3 . This sample was analyzed by alpha spectrometry which indicated the activity was radon-thoron and their respective daughters. Investigation failed to indicate a source for this activity.

During the last part of March, increases in ^{131}I concentrations were noted at the 300 and 700 Areas due to a temporary increase in the emission rate from laboratory buildings.

^{131}I concentrations increased during March and April at 100-F, 100-B, and 200-East East Center due to temporary increases in the emission rate of ^{131}I from a chemical separations facility.

May-June - The average beta concentration at most sampling stations increased in May with the spring influx of fallout, and continued higher than normal through June.

During the latter part of May, the total beta activity at 100-K reached 1.6 pCi/m^3 . A gamma spectrometer analysis of the 100-K air filter indicated $^{144}\text{Ce-Pr} = 0.41 \text{ pCi/m}^3$, $^{106}\text{Ru-Rh} = 0.47 \text{ pCi/m}^3$, and $^{95}\text{Zr-Nb} = 0.16 \text{ pCi/m}^3$. A beta decay curve on this same filter indicated two primary decaying isotopes; one with a half life of about 2.5

V. THE ATMOSPHERE (Continued)

A. Iodine-131 and Total Beta Activity (Continued)


days, corresponding closely to radon-thoron daughters, the other with a half life of about 52 days, corresponding closely to 50.4 day ^{89}Sr . Subsequent separation for ^{32}P gave a result of 0.027 pCi/m³.

The average ^{131}I and total beta concentrations for the first half of 1968 are shown below in Table 6, and for comparison the averages for 1965, 1966, and 1967 are also shown.

TABLE 6

ATMOSPHERIC ^{131}I AND TOTAL BETA CONCENTRATIONS - AVERAGES
(Results in pCi/m³)

<u>Location</u>	<u>Total Beta</u>				<u>^{131}I</u>			
	<u>1/2 1968</u>	<u>1967</u>	<u>1966</u>	<u>1965</u>	<u>1/2 1968</u>	<u>1967</u>	<u>1966</u>	<u>1965</u>
100 Areas	0.40	0.34	0.29	0.44	0.02	0.02	0.20	<0.03
200 Areas	0.31	0.41	0.58	0.80	0.02	0.09	0.10	0.14
Other On-Plant Locations	0.25	0.26	0.24	0.34	0.03	0.04	0.20	0.05




AIR SAMPLING STATIONS
● Sampling Location




Scale in Miles


IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE



IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE




IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE




IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE

--- Analytical Limit


200 W NORTHEAST



200 E WEST CENTER




200 E EAST CENTER




1967

1968


IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE



IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE



IODINE-131 AND TOTAL BETA ACTIVITY IN THE ATMOSPHERE



V. THE ATMOSPHERE (Continued)

B. Total Alpha Activity

Eleven of the 19 weekly filters which collect beta-gamma emitting radionuclides are also analyzed for alpha activity, with most of the sampling sites located in the 200 Areas. These data are presented in Figures 16-19.

For most sampling locations, results were generally at or below the analytical limit of 0.01 pCi/m³ during the first half of 1968. However, in early March the alpha concentration reached 0.913 pCi/m³ at 200-West Redox. Analysis by alpha spectrometry indicated activity was radon-thoron. An isolated temporary increase in total alpha concentration at 100-F occurred during the first week of March.

During the first part of June an alpha concentration of 0.05 pCi/m³ was reached on the 300 Area filter. Subsequent analysis by alpha spectrometry yielded the following results: ²³²Th - 36%, ²³⁸U - 20%, ²³⁴⁻²³⁵U - 42%, and ²³⁹Pu - 1.8%.

The six month average total alpha concentrations in the atmosphere for 1968 are presented in Table 7. For comparison, averages for 1965, 1966, and 1967 are also shown.


TABLE 7

ATMOSPHERIC TOTAL ALPHA CONCENTRATIONS - AVERAGES


(Results in pCi/m³)

<u>Location</u>	<u>1/2 1968</u>	<u>1967</u>	<u>1966</u>	<u>1965</u>
100 Areas	.01	.01	.01	<0.02
200 Areas	.01	.01	.01	<0.03
300 Area	.01	.02	.02	0.08
700 Area	.01	.02	.02	<0.02


TOTAL ALPHA ACTIVITY IN THE ATMOSPHERE



TOTAL ALPHA ACTIVITY IN THE ATMOSPHERE



TOTAL ALPHA ACTIVITY IN THE ATMOSPHERE



VI. RADIATION SURVEYS

A. Surface Contamination

1. Hanford Roads

Hanford roads are routinely surveyed (see Map 4) with a bio-plastic scintillation detector which is attached to the front end of a truck and is positioned about two feet above the surface of the road. The minimum level of contamination that can be detected by the road monitor corresponds to a portable GM meter response of approximately 1000 c/m per probe area. The routes between the 300 Area and the 300 Areas were surveyed bi-weekly during the report period; the remainder were surveyed monthly.

Routine surveys detected a total of ten instances of particulate contaminants during the first half of 1968, as compared to a total of seven such instances during all of 1967. Most of the increased road deposition appeared to be associated with radioactive material transport in the vicinity of the 100 Areas.

A month-by-month summary of significant findings follows:

January - Two radioactive particles were detected on Rt. 3 between 200 East and West Areas. The one with the highest reading of 100,000 c/m (GM) was recovered; the activity was found to be primarily ^{137}Cs .

February - A special survey of the road between 100-N and 100-D Areas was made on February 1 on notification that contaminated liquid dripping was suspected during waste hauling operations. A total of eight particles, including a small piece of metal, were found along the route, but no apparent liquid drops. The maximum radiation level was 150 mrads/hr (S-Juno). Laboratory analysis of the more active particles showed the activity to be a mixture of fission products and ^{60}Co . Additional surveys of road shoulders and nearby ground control plots gave negative results.

March - A maximum reading of 15,000 c/m (GM) was obtained on a particle in gravel along Rt. 2N east of 100-H Area.

Contaminated weeds at the southwest fence corner of 200 West Area read 400 c/m.

May - A particle detected along Rt. 2N at White Bluffs, reading 100,000 c/m, had ^{60}Co as the predominant radioisotope.

Two additional particles between 100-D and 100-N Area access roads were found, with the higher reading 100,000 c/m.

Numerous particulate materials associated with weeds and other organic debris were found along the dirt road south of the 200 B-C Crib Area. Several of these particles had GM readings greater than 100,000 c/m. Follow-up surveys of nearby control plots and other roads in the vicinity showed no contamination. Although more

VI. RADIATION SURVEYS (Continued)


A. Surface Contamination (Continued)

1. Hanford Roads (Continued)

numerous than previously reported, the contamination levels and extent of the spread of this material from the 200 B-C Crib Area is not believed to be significantly different from a year ago.

June - Radiation levels of 200 to 400 c/m were found on sands and gravels used for fill along George Washington Way in North Richland. Radioanalysis showed the activity to be predominantly ^{65}Zn , ^{54}Mn , and ^{137}Cs . These levels and the radionuclide spectrum are typical of the river shoreline in the vicinity of Richland. On subsequent surveys no radiation was detected.

Radioactive weeds inside the fence at the northeast corner of 200 East Area were detected, with a maximum radiation level of 150 mrad/hr (June).



Hanford Road Surveys
Survey Location


0 2 4 8
Scale in Miles

VI. RADIATION SURVEYS (Continued)

A. Surface Contamination (Continued)

2. Control Plots

Forty-six land areas, called control plots, are located within the Hanford boundaries (see Map 5). These plots, measuring 10' by 10', are periodically surveyed with a GM meter in order to detect any deposition of radioactive material. During the period covered by this report, no radioactivity was detected on the routine control plots.



CONTROL PLOT LOCATIONS
● Location

0 2 4 8

Scale in Miles

PROSSER
10 YAKIMA 47 MI.

VI. RADIATION SURVEYS (Continued)

A. Surface Contamination (Continued)

3. Waste Disposal Sites

Retired waste burial grounds are inspected periodically for general condition and evidence of disturbance. The locations of such sites outside plant areas are shown in Map 6. During the first half of 1968, inspections were made of the burial sites listed below with the findings. Where unsatisfactory conditions were found, the appropriate custodian was notified.

January


- P-11 - Good condition, no radioactivity detected.
- 100-F - Gate chain was down. This site appeared to be still active, as many waste containers were exposed within an open trench.
- 100-H - No signs posted, no radioactive materials detected.

April

- P-11 - Several small animal holes were observed, but no radioactivity was detected on the spoil from these holes.
- White Bluffs (Construction) - Good condition, no radioactivity detected.
- 100-H - No radioactive materials detected.

June

- Wye - A small pile of lead bricks with contamination levels to 3000 c/m (GM) was observed. Retirement of this site was not yet complete.
- 300 Area North - Exposed vertical piping was observed with radiation levels to 75,000 c/m (GM). Several radioactive weeds were also found with readings to 10,000 c/m. A bucket on the surface read 8000 c/m.



UNDERGROUND AND SURFACE RADIOACTIVITY

● Retired Burial Ground

VI. RADIATION SURVEYS (Continued)

A. Surface Contamination (Continued)

4. Aerial Surveys

No aerial survey flights were made during the period January-June 1968.

VI. RADIATION SURVEYS (Continued)

B. External Radiation Exposure Rates

1. Exposure Rates on-Plant

External radiation exposure rates (in units of mR/day) were determined from pencil-type gamma dosimeters located within buildings designated "614" (Map 7). The average results for the first half of 1968 are presented in Table 8 below.


Measured exposure rates were about the same in 1968 as in 1967 at most locations, although significantly higher at 100-K and 100-N Areas. Generally, the peak exposure rate during the first half of 1968 occurred during the last half of January and the first half of February (Figures 20-26), and was primarily due to regional fallout from weapons testing, although augmented by plant releases. A second peak occurred at most stations during the last half of May and the month of June which correlated with an increased influx of fallout at that time.

TABLE 8

AVERAGE EXPOSURE RATES
(Results in mR/day)

	<u>Location</u>	<u>January-June</u> <u>1968</u>	<u>January-December</u> <u>1967</u>
100 Areas	Midway	0.57	0.58
	100-B	0.82	0.72
	100-K	2.13	1.5
	100-N	1.38	0.83
	100-D	0.73	0.56
	100-F	0.68	0.39
	Hanford	0.92	0.59
200 West Area	Redox	0.56	0.59
	West Center	0.41	0.43
	East Center	0.41	0.44
200 East Area	West Center	0.52	0.37
	East Center	0.50	0.55
	Southeast	0.46	0.39
Other	Rattlesnake Springs	0.40	0.45
	Emergency Relocation Center	0.40	0.38
	*Wye Barricade	0.62	0.58
	Rt. 10 Mile 1.6	0.38	0.35
	300 Area	0.55	0.48
	700 Area	0.31	0.59

* Data for January-April only.




ON-PLANT EXTERNAL RADIATION
EXPOSURE RATE MEASUREMENT POINTS


● Measurement Point

0 2 4 8
Scale in Miles


EXTERNAL RADIATION ON PLANT




EXTERNAL RADIATION ON PLANT




EXTERNAL RADIATION ON PLANT






EXTERNAL RADIATION ON PLANT




EXTERNAL RADIATION ON PLANT



EXTERNAL RADIATION ON PLANT



EXTERNAL RADIATION ON PLANT



VI. RADIATION SURVEYS (Continued)

B. External Radiation Exposure Rates (Continued)

2. Exposure Rates at the Columbia River Shoreline

Shoreline exposure rates (Map 8 and Figures 27 and 28) are measured weekly at 3 feet above the ground with a 40-liter ionization chamber whose response is interpreted in terms of $\mu\text{R/hr}$ (radium gamma calibration). This approximates the exposure rate to the gonads of a person standing on the riverbank. Additional monthly shoreline surveys, covering the reach of the river from the reactor outfalls to Richland, include both the exposure rate at 3 feet and the levels of surface contamination as measured with a portable GM meter. The data are presented in Table 9.

In addition to the effect of the annual river flow cycle, seasonal increases in shoreline exposure rates are normally expected in March and April when increased concentrations in the Columbia River of stable ^{55}Mn due to spring run-off result in increased concentrations of ^{56}Mn .

During the first half of 1968, shoreline exposure rates and particulate frequency at locations below the plant boundary continued to show increases compared to 1966-1967. These increases were observed during the latter half of 1967 and were attributed to a combination of reactor area events and river behavior. The effects of rapidly changing river flow rates were also seen in late June when particulate contamination was detected on a previously-flooded area of the plant shore near PRTR. As usual, the exposure rates decreased in late May when the river flow rate increased.

Unusual results during the period included the following.

January - A shoreline exposure rate of 1500 $\mu\text{R/hr}$ was measured below 100-K. Exposure rates of 50% higher than in 1967 were observed downstream as far as Ringold.

February - Particles of 60,000 c/m were found at both Ringold and the Power Line Crossing during low water following large river fluctuations. Analysis showed predominantly ^{46}Sc , but a number of other radionuclides were detectable.

March - A particle reading 50,000 c/m was found at the Hanford Far Shore location. Analysis showed ^{65}Zn predominating with ^{46}Sc , $^{144}\text{Ce-Pr}$, ^{51}Cr , and ^{155}Eu also present.

A maximum reading of 4,000 c/m was obtained on mud at the Richland Pump Plant.


May - Several particles with a maximum reading of 75,000 c/m were detected at Ringold and the Power Line Crossing. Patches of foam supporting mixed debris read 25,000 c/m at 100-F Area and 9,000 c/m at Sacajawea Park.

VI. RADIATION SURVEYS (Continued)


B. External Radiation Exposure Rates (Continued)

2. Exposure Rates at the Columbia River Shoreline (Continued)

June - Numerous particles with a maximum of 8,000 c/m were detected on a previously flooded area near the PRTR pump intake. After re-flooding and re-exposure, most of the particulate material was gone, and a maximum reading of 4,000 c/m was obtained on the remaining debris.



EXTERNAL RADIATION AT THE COLUMBIA RIVER SHORELINE



EXTERNAL RADIATION AT THE COLUMBIA RIVER SHORELINE




TABLE 9
MONTHLY RIVER SHORELINE SURVEYS - JANUARY-JUNE, 1968
(Maximum Readings)

Location	January 17 μR/hr c/m @ @ 3ft. Surf.	February 2 μR/hr c/m @ @ 3ft. Surf.	March 21 μR/hr c/m @ @ 3ft. Surf.	May 3 μR/hr c/m @ @ 3ft. Surf.	May 28 μR/hr c/m @ @ 3ft. Surf.	June 27 μR/hr c/m @ @ 3ft. Surf.
Above 181-KW-Plant Shore	90	500	64	300	45	56
Below 181-KW-Plant Shore	140	800	180	1000	70	67
Below 181-KW-Far Shore	26	250	22	100	40	14
Above 181-NE-Plant Shore	130	400	100	700	36	20
Below 181-NE-Plant Shore	130	2000	100	600	28	26
Above 181-D-Far Shore	20	250	22	100	24	18
D Island-Far Shore	300	2500	240	3000	54	46
Below 181-D-Plant Shore	125	800	72	400	55	28
Below 181-D-Far Shore	36	250	28	100	23	15
Below 181-H-Plant Shore	260	2500	98	600	40	36
White Bluffs Ferry-Plant Shore	180	800	74	400	30	23
White Bluffs Ferry-Far Shore	86	600	38	150	28	15
Hanford-Plant Shore	175	1000	76	500	58	23
Hanford-Far Shore	75	500	81	300	27	22
Ringold-Far Shore	76	600	42	300	26	18
Powerline-Plant Shore	200	1000	100	600	27	37
Powerline-Far Shore	170	600	74	500	22	23
Byer's Landing-Far Shore	56	250	65	250	56	38
PRTR-Plant Shore	86	600	82	1000	32	26
Richland Ferry-Plant Shore	42	400	50	1500	42	24


Notes: Measurements reported in μR/hr are taken with a 40-liter ionization chamber, the center of the chamber 3 ft. above the water's edge; measurements reported in c/m are the maximum found with a GM in the immediate area of the water's edge.

VI. RADIATION SURVEYS (Continued)

B. External Radiation Exposure Rates (Continued)

3. Exposure Rates Below the Surface of the Columbia River

Exposure rates in the river were determined from pocket dosimeters contained within submerged bottles at the locations shown in Map 9. Missing data were the result of lost containers or equipment malfunctions. (See Figures 29 and 30). Average dose rates for the first half of 1968 at the 300 Area and Richland were somewhat lower than for the same period of 1967.




EXPOSURE RATE MEASUREMENT POINTS
IN THE COLUMBIA RIVER

- Sub-Surface Measurement Point

0 2 4 8
Scale in Miles

EXTERNAL RADIATION BELOW THE SURFACE OF THE COLUMBIA RIVER



EXTERNAL RADIATION BELOW THE SURFACE OF THE COLUMBIA RIVER

